

# ***rbcL* Transcript Levels in Tobacco Plastids Are Independent of Light: Reduced Dark Transcription Rate Is Compensated by Increased mRNA Stability**

Takashi Shiina,<sup>1</sup> Lori Allison,<sup>2</sup> and Pal Maliga<sup>3</sup>

Waksman Institute, Rutgers, The State University of New Jersey, 190 Frelinghuysen Road, Piscataway, New Jersey 08854-8020

The plastid *rbcL* gene, encoding the large subunit of ribulose-1,5-bisphosphate carboxylase, in higher plants is transcribed from a  $\sigma^{70}$  promoter by the eubacterial-type RNA polymerase. To identify regulatory elements outside of the *rbcL* –10/–35 promoter core, we constructed transplastomic tobacco plants with *uidA* reporter genes expressed from *rbcL* promoter derivatives. Promoter activity was characterized by measuring steady state levels of *uidA* mRNA on RNA gel blots and by measuring promoter strength in run-on transcription assays. We report here that the *rbcL* core promoter is sufficient to obtain wild-type rates of transcription. Furthermore, the rates of transcription were up to 10-fold higher in light-grown leaves than in dark-adapted plants. Although the rates of transcription were lower in the dark, *rbcL* mRNA accumulated to similar levels in light-grown and dark-adapted leaves. Accumulation of *uidA* mRNA from most *rbcL* promoter deletion derivatives directly reflected the relative rates of transcription: high in the light-grown and low in the dark-adapted leaves. However, *uidA* mRNA accumulated to high levels in a light-independent fashion as long as a segment encoding a stem-loop structure in the 5' untranslated region was included in the promoter construct. This finding indicates that lower rates of *rbcL* transcription in the dark are compensated by increased mRNA stability.

## **INTRODUCTION**

Plastid genes are transcribed by two RNA polymerases, one of which is related to the eubacterial  $\alpha_2\beta\beta'$  RNA polymerase (Igloi and Kössel, 1992; Grissem and Tonkyn, 1993). The  $\alpha_2\beta\beta'$  homologs are encoded by the plastid genome; therefore, this RNA polymerase is referred to as the plastid-encoded plastid RNA polymerase (PEP). PEP promoter recognition is probably dependent on  $\sigma$ -like factors (Tiller et al., 1991; Tiller and Link, 1993a), which are encoded in the nucleus (Liu and Troxler, 1996; Allison, 1997; Isono et al., 1997a; Tanaka et al., 1997; Tozawa et al., 1998). PEP promoters are similar to the eubacterial  $\sigma^{70}$ -type promoters: core promoters comprise two conserved blocks of hexameric sequences (consensus; TTGACA and TATAAT) corresponding to the eubacterial –35 and –10 promoter elements. The hexamers are spaced 17 to 19 nucleotides apart; transcription initiates five to seven nucleotides downstream of the –10 box sequence. Accumulation of transcripts at a high level from poorly conserved PEP promoters depends on up-

stream activators such as the light-responsive promoter of the *psbD* gene encoding the D2 photosystem II subunit protein (Allison and Maliga, 1995; Kim and Mullet, 1995; To et al., 1996; Satoh et al., 1997). The second plastid RNA polymerase is nuclear encoded (NEP) and is likely to resemble a phage-type single-subunit enzyme related to the mitochondrial RNA polymerase (Lerbs-Mache, 1993; Hedtke et al., 1997). NEP promoters have a short, conserved sequence around the transcription initiation site (Hajdukiewicz et al., 1997; Kapoor et al., 1997; Hübschmann and Börner, 1998; Silhavy and Maliga, 1998a). PEP transcribes photosystem I and II genes, and NEP transcribes *accD* and the *rpoB* operon, whereas both polymerases transcribe the remaining genetic system and metabolic genes (reviewed in Maliga, 1998).

The plastid *rbcL* gene encodes the large subunit of ribulose-1,5-bisphosphate carboxylase (EC 4.1.1.39). The *rbcL* gene is transcribed by PEP from a single promoter, which has been mapped in a number of species, including tobacco (Shinozaki and Sugiura, 1982), maize, spinach, pea (Mullet et al., 1985), barley (Reinbothe et al., 1993), and Arabidopsis (Isono et al., 1997b). The tobacco *rbcL* gene has a PEP promoter with relatively well-conserved –35 and –10 elements (TTGCGC and TACAAT) and canonical spacing (18 nucleotides). In vitro studies have confirmed the importance of the –35 and –10 box spacing and sequence for *rbcL*

<sup>1</sup>Current address: Graduate School of Human and Environmental Studies, Kyoto University, Yoshida-nihonmatsu-mati, Sakyo-ku, Kyoto 606-8501, Japan.

<sup>2</sup>Current address: N258 Beadle Center, University of Nebraska, Lincoln, NE 68588-0664.

<sup>3</sup>To whom correspondence should be addressed. E-mail maliga@waksman.rutgers.edu; fax 732-445-5735.

promoter strength (Gruissem and Zurawski, 1985; Hanley-Bowdoin et al., 1985).

In vitro plastid transcription assays do not always reproduce the in vivo expression pattern (Wada et al., 1994). Therefore, we decided to test the role of sequences outside of the *rbcl* promoter core by expression of *uidA* reporter genes from *rbcl* promoter derivatives in vivo by using a transgenic approach (Svab and Maliga, 1993).

Sequences positioned between nucleotides  $-16$  and  $-102$  relative to the *rbcl* transcription initiation site are proposed to function in maize as a binding site for the chloroplast DNA binding factor 1 (CDF1) sequence-specific transcription factor. Segments of the CDF1 binding site are conserved between maize, pea, spinach, and tobacco (Lam et al., 1988). The region corresponding to the putative CDF1 binding site in tobacco is between nucleotides  $-16$  and  $-96$  with three conserved regions: promoter-distal CDF1 region I contains seven nucleotides of the maize GTATTTAG repeat; CDF1 region II corresponds to two A-rich sequences termed A boxes, whereas most of CDF1 region III is within the promoter core (underlined in Figure 1). We prepared a set of promoter deletion derivatives to test the role of the upstream elements in *rbcl* promoter function. DNA sequences downstream of the transcription initiation site were also tested for


contribution to promoter activity. Although not yet confirmed for a higher plant, such *rbcl* promoter elements were found in the unicellular alga *Chlamydomonas* (Klein et al., 1994).

We report here that the *rbcl* core promoter is sufficient to obtain wild-type rates of transcription of the chimeric *uidA* genes. Therefore, no sequences outside of the promoter core significantly contribute to *rbcl* promoter function. Furthermore, the rates of *rbcl* transcription were significantly reduced in dark-adapted plants. Reduced rates of transcription resulted in lower steady state *uidA* mRNA levels in the dark, unless the 5' segment of the *rbcl* 5' untranslated region (5' UTR) was included in the chimeric construct. This finding indicates that accumulation of *rbcl* mRNA in a light-independent manner is due to stabilization of the mRNA via its 5' UTR compensating for reduced rates of transcription in the dark.

## RESULTS

### The *rbcl* Promoter Derivatives

A series of promoter deletion derivatives was constructed to test the role of sequences outside of the *rbcl* core for their



**Figure 1.** Chimeric *uidA* Genes to Test *rbcl* Promoter Activity.

*Prbcl* indicates the position of the *rbcl* promoter deletion derivatives; *uidA* encodes the  $\beta$ -glucuronidase reporter enzyme; *Trps16* refers to the 3' UTR of the *rps16* ribosomal protein gene. The *rbcl* promoter sequence, with the transcription initiation site (horizontal arrow), the  $-10/-35$  promoter elements, and conserved CDF1 regions (underlined) are shown below the chimeric gene. The *rbcl* promoter derivatives are identified by the pWW plasmid number.

contribution to promoter activity (Figure 1). The longest *rbcl* promoter in the 5' deletion series includes CDF1 region I (−96/−89), the two A-rich A boxes (A box 1, −66/−54; A box 2, −51/−40), and nine nucleotides of the 5' UTR in plasmid pWW11. In the shorter derivatives, one or more of these sequences have been systematically removed (pWW12, pWW14, pWW15, and pWW16; Figure 1).

Promoters in plasmids pWW12, pWW13, and pIK100 were designed to test for functional promoter elements downstream of the −35/−10 promoter core. These plasmids contain nine, 122, and 180 nucleotides downstream of the transcription initiation site, respectively (Figure 1). Note that the promoter driving *uidA* expression in plasmid pWW12 is shown in both the 5' and the 3' deletion series.


The *rbcl* promoter derivatives were cloned upstream of a *uidA* construct consisting of a ribosome binding site; *uidA*, encoding  $\beta$ -glucuronidase; and *Trps16*, the 3' UTR of the *rps16* ribosomal protein gene to stabilize the mRNA. The chimeric *uidA* genes were cloned into the pPRV112A plastid transformation vector (Zoubenko et al., 1994) and introduced into the inverted repeat region of the tobacco plastid genome (Figure 2).

#### Accumulation of mRNA to Wild-Type Levels in the Dark Depends on Sequences Downstream of the $\sigma^{70}$ -Type Core Promoter

Steady state levels of the *rbcl* mRNA are similar in light-grown and dark-adapted tobacco plants (see below). Preliminary data indicated light-dependent accumulation of reporter mRNAs in some of the transgenic plants. Therefore, we studied mRNA accumulation in the leaves of dark-adapted plants (96 hr; DD plants) versus dark-adapted plants that had been illuminated for 24 hr (DL plants). Steady state mRNA levels for *rbcl* and the chimeric *uidA* genes are similar in DL plants and in plants grown in cycling light (16 hr of light and 8 hr of dark; data not shown).

In the transgenic plants, we probed for *rbcl* mRNA to examine mRNA accumulation from the native *rbcl* promoter and *uidA* mRNA to examine mRNA accumulation from the chimeric *rbcl* promoters. As a control, the 16S rRNA was used to detect RNA accumulation from the native rRNA operon promoter because accumulation of mRNA from the  $\sigma^{70}$ -type *Prrn-114* promoter is not dependent on light (Allison and Maliga, 1995). To control for equal loading, we probed the blots for the cytoplasmic 25S rRNA.

Accumulation of the *uidA* mRNA in the illuminated (DL) plants was comparable in all of the transgenic lines. Accumulation of *uidA* mRNA in the dark-adapted (DD) samples was lower than in the illuminated (DL) samples, unless the 5' segment of the *rbcl* 5' UTR between nucleotides +9 and +122 was incorporated in the chimeric promoters. The importance of this segment can be seen by comparing levels of mRNA in dark-adapted plants transformed with plasmids pWW12 and pWW13 (Figure 3).



**Figure 2.** Map of the Tobacco Plastid Genome with the Selectable Marker *aadA* and Chimeric *PrbcL::uidA::Trps16* Gene.


The transgenes have been inserted into the *trnV-rps7/12* intergenic region. The 16S rDNA and *rps7/12* are plastid genes (Shinozaki et al., 1986). Wavy horizontal lines represent transcripts detected by the *uidA* probe in the pWW transgenic series and in the pIK100 transgenic plant. The asterisk marks the readthrough transcript. Arrows indicate the orientation of plastid genes. Hatched and stippled bars flanking *aadA* and *uidA* symbolize 5' and 3' regulatory regions.

There were two alternative explanations for high levels of *uidA* mRNA in the dark-adapted pWW13 and pIK100 plants. The pWW13 promoter fragment may have contained a promoter element downstream of the  $\sigma^{70}$  core that is responsible for maintaining high (illuminated) levels of transcription in the dark. Alternatively, the segment between positions +9 and +122 directed stabilization of the *uidA* mRNA in dark-adapted leaves.

#### A Segment of the *rbcl* 5' UTR Protects the pWW13 *uidA* mRNA from Degradation

The two alternatives could be distinguished by measuring rates of transcription. Higher and lower rates of transcription correlating with higher and lower levels of steady state mRNAs in the dark-adapted plants would indicate a promoter element that enhances *rbcl* transcription in the dark. If, however, transcription rates from the chimeric promoters are higher in illuminated (DL) plants and lower in dark-adapted (DD) plants independent of the levels of mRNA, high steady state levels of *uidA* mRNAs in the pWW13 and pIK100 plants should be due to stabilization of the mRNAs in the dark.

In organello rates of transcription are determined by run-on transcription assays. First, chloroplasts are isolated while preserving plastid transcriptional elongation complexes. Transcriptional elongation is then continued in the presence of radioactive nucleotides. Radioactivity incorporated into specific mRNAs is determined by hybridization to immobilized DNA probes (Deng and Gruissem, 1995). The results of such a slot blot hybridization with the transgenic plants are



**Figure 3.** Accumulation of mRNAs from the Chimeric *uidA* Genes.

RNA samples were derived from dark-adapted (96 hr) plants (DD lanes) and dark-adapted (72 hr) and illuminated (24 hr) plants (DL lanes). The amount of total cellular RNA per lane was 3  $\mu$ g for *rbcL* and *uidA* and 1  $\mu$ g for the 16S rRNA. An asterisk marks a readthrough dicistronic (*uidA-aadA*) transcript in the pIK100 plastids. WT, wild type.

shown in Figure 4. Relative activities, normalized for the probe size, are given in Figure 5. The amount of incorporated UTP in plastid RNA in the dark-adapted (DD) plants was in the range of 0.33 to 1.49 pmol  $5 \times 10^6$  plastids $^{-1}$  5 min $^{-1}$ , whereas in the illuminated (DL) plants, it was in the range of 1.04 to 5.24 pmol  $5 \times 10^6$  plastids $^{-1}$  5 min $^{-1}$ . These values are similar to those found in 8-day-old barley seedlings (Figure 1 in Klein and Mullet, 1990).

The rate of *rbcL* transcription relative to the 16S rDNA is  $\sim 40\%$  (Figure 5). This is also true of the chimeric *rbcL* promoters, indicating that all sequences relevant for transcription initiation from the *rbcL* promoter are present, even in the smallest fragment in the pWW16 construct. For both the native *rbcL* and the control 16S rDNA genes, the rates of transcription were relatively high in the illuminated (DL) sample and dropped dramatically (four- to 10-fold) in the dark-adapted (DD) sample. The pWW13 and pIK100 plants were no exception: transcription of the *uidA* gene was significantly reduced in the dark-adapted leaves, indicating that higher *uidA* mRNA levels in the dark are due to stabilization of the mRNA.


## DISCUSSION

### The *rbcL* $\alpha^{70}$ -Type Promoter Core Is Sufficient for Normal Rates of Transcription in Vivo

Data presented in Figure 5 indicate that no sequence outside of the *rbcL* promoter core is necessary for full promoter activity. This conclusion is based on finding close to wild-type rates of *rbcL* transcription in all transgenic plants, including the pWW16 line, which carries the smallest (–35 to +9) *rbcL* promoter derivative, containing only a slightly larger fragment than the promoter core (–35 to –5).


It should be noted that the wild-type plants carry a single *rbcL* promoter per plastid genome, whereas the transgenic plants carry three: the wild-type *rbcL* promoter in the large single-copy region and two copies driving *uidA* expression in the inverted repeat region. Interestingly, introduction of two additional *rbcL* promoter copies does not effect *rbcL* promoter activity.

CDF1 binding in maize was more evident in bundle sheath chloroplasts than in mesophyll cell chloroplasts. Because CO<sub>2</sub> fixation by ribulose-1,5-bisphosphate carboxylase occurs in bundle sheath cells, CDF1 was proposed to be a cell type-specific regulator of *rbcL* expression (Lam et al., 1988).



**Figure 4.** Slot Blot Analysis to Measure Rates of Plastid Gene Transcription.

RNA samples derived from dark-adapted (96 hr) plants (DD lanes) and dark-adapted (72 hr) and illuminated (24 hr) plants (DL lanes). Rates of transcription were measured for the 16S rRNA, *rbcL*, and chimeric *uidA* genes expressed from the *rbcL* promoter derivatives. Plasmid pBluescript II KS+ (pBS) DNA was used as a control for nonspecific hybridization of the run-on transcripts. The nonlabeled DNA (0.5 pmol per slot) was probed with  $^{32}$ P-labeled RNA from the lysed transcription assay of  $5 \times 10^6$  chloroplasts. Data shown on this representative blot may be slightly different from the averages of multiple experiments shown in Figure 5. WT, wild type.



**Figure 5.** Relative Rates of Transcription of the 16S rDNA, *rbcL*, and Chimeric *uidA* Genes.

Gene-specific activities were calculated as UTP incorporated (picomoles) per  $10^6$  plastids per 1-kb length of DNA probe during a 5-min assay. The values are given as the percentage of the 16S rDNA transcription rate in the light (100%) in arbitrary units (mean  $\pm$  SD). Under the experimental conditions used here, the actual 16S rDNA transcription rate may be underestimated, resulting in overestimation of the relative *rbcL* transcription rates. The values were calculated from four to six independent slot blot hybridization experiments, except for the wild-type plants, for which the numbers were calculated from two experiments. Abbreviations are as given in the legend to Figure 4.

This still may be the case. Given the limited sequence conservation between maize and tobacco CDF1 binding sites, and because tobacco is a  $C_3$  rather than a  $C_4$  plant, the lack of a role for the CDF1 chloroplast DNA binding factor in tobacco does not necessarily exclude a role for CDF1 sequences in  $C_4$  plastids.

In *Escherichia coli*, the AT-rich UP element is localized between nucleotides  $-40$  to  $-60$  upstream of the transcription initiation site and stimulates transcription by a factor of 30 in vivo (Ross et al., 1993). Tobacco *rbcL* CDF1 region II (A boxes) is similarly spaced relative to the *rbcL* promoter core. Sequences upstream of the  $-35$  box in the *psbD* promoter contain binding sites for a factor regulating *psbD* transcription (Allison and Maliga, 1995; Kim and Mullet, 1995; Christopher, 1996; To et al., 1996; Satoh et al., 1997). However, the two A box regions upstream of the tobacco *rbcL* promoter do not seem to play a role in regulating transcription. This is also true of sequences downstream of the promoter core that are required for full *rbcL* promoter strength in *Chlamydomonas* (Klein et al., 1994).

#### Transcription from $\sigma^{70}$ -Type Promoters Is Light Regulated in Chloroplasts

We have shown that steady state levels of *rbcL* mRNA in tobacco are comparable in dark-adapted (DD) and light-

induced (DL) leaves. Also, *uidA* mRNA accumulates to similar levels in dark-adapted (DD) and light-induced (DL) leaves from the *Prrn-114* promoter, which drives rRNA transcription in leaves (Allison and Maliga, 1995; Vera and Sugiura, 1995; Allison et al., 1996). Given light-independent accumulation of mRNAs from these  $\sigma^{70}$ -type PEP promoters, we had expected to find that transcription from these promoters is constitutive. Interestingly, this is not the case. Transcription rates in dark-adapted leaves are four to 10 times lower than in the illuminated leaves for both *rbcL* and the 16S rDNA. Enhanced rates of plastid transcription by light have been reported in several species, including spinach (Deng and Gruissem, 1987), barley, maize (Baumgartner et al., 1993), sorghum (Schrubar et al., 1990), wheat (Kawaguchi et al., 1992), pea (DuBell and Mullet, 1995), and Arabidopsis (Hoffer and Christopher, 1997). Responsiveness of plastid gene transcription to light also depends on the developmental stage of the plastid (Klein and Mullet, 1990; Baumgartner et al., 1993).

Multiple mechanisms may lead to light-induced PEP activity. Light-induced synthesis of the components of the PEP transcription machinery may be one such mechanism. Candidates for this type of regulation are the PEP  $\sigma$  factor genes, whose transcription is induced by light (Liu and Troxler, 1996; Isono et al., 1997a; Tanaka et al., 1997; Tozawa et al., 1998). Conversion of the eubacterial-type PEP core (B form) into the more complex A form during light-induced development


may also contribute to light-induced PEP activity (Pfannschmidt and Link, 1994, 1997). An additional mechanism could be reversible phosphorylation of the  $\sigma$  factors shown to regulate rates of transcription in vitro (Tiller and Link, 1993b) and in vivo (Christopher et al., 1997). The in vivo role of phosphorylation in regulating plastid gene transcription seems to be corroborated by the purification of a PEP-associated protein kinase, which phosphorylates chloroplast  $\sigma$ -like factors in vitro (Baginsky et al., 1997).

### The 5' UTR Is Responsible for Stabilizing the *rbcl* mRNA in the Dark

The experimental observations of light-dependent *rbcl* transcription and similar steady state levels of *rbcl* mRNA in dark-adapted (DD) and illuminated (DL) leaves suggest stabilization of mRNA as a compensating mechanism for the lower rates of transcription in the dark. The in vivo study of *uidA* expression from *rbcl* promoter derivatives points to the 5' UTR as the source of the stabilizing sequences. In the plastids of higher plants, studies of mRNA stability thus far have been limited to the role of the 3' UTR. The 3' UTR frequently contains a short stem-loop structure associated with a high molecular weight complex. This complex contains a 3' to 5' exonuclease (PNPase, or 100-kD polynucleotide phosphorylase), two endoribonucleases, and at least four additional proteins involved in 3' end formation and degradation of the mRNA (Stern and Gruissem, 1987; Hayes et al., 1996; Yang et al., 1996). A putative short stem-loop structure is also present in the *rbcl* 5' UTR. This is disrupted by the deletion in the pWW12 promoter derivative but is present in the pWW13 and pK100 promoters (Figure 6). The segment of the *rbcl* 5' UTR containing this stem-loop structure is highly conserved during evolution, suggesting function in vivo (Figure 6).

The tobacco *rbcl* mRNA has an RNA processing site in the 5' UTR between nucleotide positions +123 and +124, which are marked in Figures 1 and 6 (Hanley-Bowdoin et al., 1985; Mullet et al., 1985; Reinbothe et al., 1993). This RNA processing site is absent in all *rbcl* promoter derivatives except pK100 (Figure 1). The chimeric mRNA is unstable in dark-adapted (DD) pWW12 plants, whereas it is stabilized in pWW13 plants (Figure 3). Both sets of transgenic plants lack the RNA processing site. Therefore, stabilization of chimeric mRNAs is independent of RNA processing at this site.

We report here regulation of plastid mRNA stability via the mRNA 5' UTR. To our knowledge, no such information has been published for a higher plant. However, there are several well-documented cases for regulation of mRNA stability via the 5' UTR in the unicellular alga *Chlamydomonas*. The examples include studies of nuclear mutations that target the 5' UTR and thereby destabilize the *psbD* mRNA (reviewed in Mayfield et al., 1995; Rochaix, 1995). A transgenic approach was used to confirm the role of the *rbcl* (Salvador et al., 1993) and *petD* (Sakamoto et al., 1993, 1994) 5' UTRs and of the *atpB* mRNA



**Figure 6.** DNA Sequence Alignment of Regions Containing the *rbcl* 5' UTR in Higher Plants.

Nt, tobacco (Shinozaki and Sugiura, 1982); At, Arabidopsis (Isono et al., 1997b); Ps, pea (Zurawski et al., 1986); Os, rice (Silhavy and Maliga, 1998b); Hv, barley (Reinbothe et al., 1993); and Zm, maize (Mullet et al., 1985). Nucleotides conserved in all six species are boxed; those conserved in five of the six species are marked by asterisks under the alignment. The *rbcl* translation initiation codon (ATG) is shown in boldface. Positions of the -10 and -35 promoter elements, the Shine-Dalgarno (SD) sequence, the processing site, and the 3' end of promoter derivatives (filled triangles) are marked above the sequence; inverted repeats in tobacco are marked by horizontal arrows below the sequence. The size of the repeated region is slightly different in the other species. Dashes were introduced to optimize alignment.

3' UTR (Stern et al., 1991; Stern and Kindle, 1993) in mRNA maturation and turnover. The role of the short stem-loop structure, and possibly of other RNA *cis* sequences, in stabilizing the higher plant *rbcL* mRNAs in the dark remains to be determined.

## METHODS

### Plasmid Construction

A series of 5' deletions of the upstream sequence of the *rbcL* transcription initiation site was amplified using pK36 containing the *rbcL* promoter as a template. Plasmid pK36 is a pBluescript KS+ (Stratagene, La Jolla, CA) vector derivative carrying an *SpeI*-*SacII* plastid DNA fragment spanning nucleotides 56,389 to 57,750 in the plastid genome (Shinozaki et al., 1986). The *rbcL* promoter derivatives were amplified as *SacI* (at the 5') and *EcoRI* (at the 3') fragments with the following primers: pWW11, 5'-GGGAGCTCGGGGAAGTTCTTATTA-3' (5' end point -110) and 5'-GGAATTCAAATACATCAT-TATTG-3' (3' end point at +9); pWW12, 5'-GGGAGCTCGTCGACTAGTCAGGTATTTTC-3' (5' end point -86) and 5'-GGAATTCAAATACATCAT-TATTG-3' (3' end point at +9); pWW13, 5'-GGGAGCTCGTCGACTAGTCAGGTATTTTC-3' (5' end point -86) and 5'-CGAATTCGGTGAATGA-3' (3' end point at +122). The fragments obtained by amplification with these primers were digested with *EcoRI* and *SacI* and ligated into *SacI*-*EcoRI*-digested plasmid pLAA25A (Zoubenko et al., 1994) to obtain plasmids pWW12, pWW13, pWW14, pWW15, and pWW16. Plasmid pLAA25A is a pRRV112A plastid transformation vector derivative with a selectable spectinomycin resistance (*aadA*) gene and a promoterless *uidA* coding region with (1) a synthetic ribosome binding site and (2) the *rps16* ribosomal protein gene 3' untranslated region (UTR; *Trps16*; Zoubenko et al., 1994). Cloning of *rbcL* promoter fragments upstream of the ribosome binding site created the pWW series of *PrbcL::rbcL(S)::uidA::Trps16* reporter genes.

Plasmid pK100 is a pPRV111B plastid transformation vector derivative (Zoubenko et al., 1994) in which the *uidA* reporter gene (*PrbcL::uidA::TrbcL*) is cloned as a *Sall*-*HindIII* fragment in the poly-cloning site in the same orientation as the selectable marker spectinomycin resistance (*aadA*) gene. *PrbcL* in plasmid pK100 is contained in a *Sall* (at the 5' end) and *NcoI* (3') fragment, including nucleotides between 57,319 and 57,586 of the plastid genome (Shinozaki et al., 1986). This region includes sequences -86/+181 relative to the transcription initiation site. The *Sall* and *NcoI* sites were created by site-directed mutagenesis using oligonucleotides 5'-GGAAATACCTGACTAGTCGACCTAAATAATAAGAACTTC-3' and 5'-GTCTCTGTTGTGGTGCCATGGATCCCTCCCTACAAC-3', respectively. The *uidA* coding segment is identical in the pWW plasmids and in plasmid pK100, and it is contained in an *NcoI*-*XbaI* fragment. *TrbcL* (*XbaI*-*HindIII* fragment) has been described previously (Staub and Maliga, 1994). The chimeric *PrbcL::uidA::TrbcL* gene was cloned in plastid vector pPRV111B (Zoubenko et al., 1994) to yield plasmid pK100. Plants

transformed with plasmid pK100 were made available by I. Kanevski (Waksman Institute).

### Plastid Transformation

Tobacco (*Nicotiana tabacum* cv Petit Havana) plants were grown aseptically on RM medium containing agar-solidified Murashige and Skoog salts (Murashige and Skoog, 1962) and sucrose (3%). Tungsten particles (1  $\mu$ m) coated with plasmid DNA were introduced into the leaves by using the DuPont PDS1000He Biolistic gun at 1100 psi (Svab and Maliga, 1993). Transgenic shoots were selected on RMOP medium containing 500 mg/L spectinomycin dihydrochloride (Svab et al., 1990). Homoplastomic transformed lines were identified by DNA gel blot analysis after two cycles of plant regeneration on the same selective spectinomycin medium. Transgenic cuttings were rooted on Murashige and Skoog agar medium (Murashige and Skoog, 1962).

### Light Treatment

Transgenic plants were rooted aseptically on solid RM media in culture boxes and maintained under a 16-hr-light and 8-hr-dark cycle for 4 to 8 weeks. For dark adaptation, plants were placed in complete darkness for 3 days. Subsequently, plants were either maintained in the dark or returned to white light (cool-white fluorescent bulbs at  $\sim 10,000$  lux) for 24 hr.

### RNA Gel Blot Analysis

Total cellular RNA was isolated from fully expanded leaves frozen in liquid nitrogen by using TRIzol (Gibco BRL), following the manufacturer's protocol. The RNA was electrophoresed on 1% agarose-formaldehyde gels and transferred to a Hybond N membrane (Amersham) by using the PosiBlot transfer apparatus (Stratagene). Blots were hybridized using the rapid hybridization buffer (Amersham) at 65°C, with  $^{32}$ P-labeled double-stranded DNA probes generated by random primed labeling (Boehringer Mannheim).

### Chloroplast Isolation and Run-on Transcription

The plants were grown aseptically on RM medium containing agar-solidified Murashige and Skoog salts and 3% sucrose in Magenta boxes. Chloroplast run-on transcription assays were performed according to Deng and Gruissem (1995). Briefly, leaves (5 to 10 g) of aseptically grown plants were homogenized in an isolation buffer (25 mM Hepes-KOH, pH 7.6, 350 mM sorbitol, and 2 mM EDTA) in a Waring blender. Intact plastids were isolated by centrifugation (10,000g) on a 10 to 80% Percoll gradient. The chloroplast number was determined by counting in a hemocytometer. Run-on transcription was performed using  $\alpha$ - $^{32}$ P-UTP (specific activity of 400 mCi/mmol) and  $10^7$  chloroplasts in the presence of 20 mg/mL heparin. After an 8-min incubation at 25°C, duplicate aliquots were spotted on DE-81 paper (Whatman, Clifton, NJ) for measurement of total UTP incorporation into RNA. Radioactive run-on transcripts were purified by phenol extraction and hybridized with nonradiolabeled 16S rDNA, *rbcL*, and *uidA* gene probes and pBS plasmid DNA (0.5 pmol each) slot blotted on nylon membranes. The probes were the 16S rDNA 1.28-kb *EcoRI*-*SphI* fragment (Shinozaki et al., 1986); *rbcL* 1.3-kb *EcoRI*-*XbaI* fragment (Shinozaki et al., 1986); *uidA* 1.87-kb *NcoI*-*XbaI*

fragment (plasmid pLAA25A; Zoubenko et al., 1994); and plasmid pBluescript II KS+ EcoRI-digested plasmid DNA (Stratagene). Slot blots were probed with lysed assays of  $5 \times 10^6$  chloroplasts. Radioactivity hybridizing with the probes was determined by measuring radioactivity in excised bands by using a liquid scintillation counter.

## ACKNOWLEDGMENTS

We thank Wei Wu for her contributions to plasmid construction and plant transformation during the early phase of the experiments and Dr. Ivan Kanevski for plasmid pIK36 and for tobacco plants transformed with plasmid pIK100. Research was supported by a grant from the National Science Foundation to P.M. Contributions by T.S. were made feasible by a grant from the Research for the Future program, Japan Society for the Promotion of Science (No. JSPS-RFTF96L00604).

Received May 20, 1998; accepted August 4, 1998.

## REFERENCES

- Allison, L.A. (1997). Multiple sigma factors provide a mechanism for nuclear control over chloroplast gene expression. *Plant Physiol.* **114** (suppl.), 65 (abstr.).
- Allison, L.A., and Maliga, P. (1995). Light-responsive and transcription-enhancing elements regulate the plastid *psbD* core promoter. *EMBO J.* **14**, 3721–3730.
- Allison, L.A., Simon, L.D., and Maliga, P. (1996). Deletion of *rpoB* reveals a second distinct transcription system in plastids of higher plants. *EMBO J.* **15**, 2802–2809.
- Baginsky, S., Tiller, K., and Link, G. (1997). Transcription factor phosphorylation by a protein kinase associated with chloroplast RNA polymerase from mustard (*Sinapis alba*). *Plant Mol. Biol.* **34**, 181–189.
- Baumgartner, B.J., Rapp, J.C., and Mullet, J.E. (1993). Plastid genes encoding the transcription/translation apparatus are differentially transcribed early in barley (*Hordeum vulgare*) chloroplast development. *Plant Physiol.* **101**, 781–791.
- Christopher, D.A. (1996). Leaf development and phytochrome modulate the activation of *psbD-psbC* transcription by high-fluence blue light in barley transcripts. *Photosynth. Res.* **47**, 239–251.
- Christopher, D.A., Xinli, L., Kim, M., and Mullet, J. (1997). Involvement of protein kinase and extraplastidic serine/threonine protein phosphatase in signaling pathways regulating plastid transcription and the *psbD* blue light responsive promoter in barley. *Plant Physiol.* **113**, 1273–1282.
- Deng, X.W., and Gruissem, W. (1987). Control of plastid gene expression during development: The limited role of transcriptional regulation. *Cell* **49**, 379–387.
- Deng, X.W., and Gruissem, W. (1995). Chloroplast run-on transcription: Determination of the transcriptional activity of chloroplast genes. In *Methods in Plant Molecular Biology*, P. Maliga, D.F. Klessig, A.R. Cashmore, W. Gruissem, and J.E. Varner, eds (Plainview, NY: Cold Spring Harbor Laboratory Press), pp. 191–207.
- DuBell, A.N., and Mullet, J.E. (1995). Differential transcription of pea chloroplast genes during light-induced leaf development. *Plant Physiol.* **109**, 105–112.
- Gruissem, W., and Tonkyn, J.C. (1993). Control mechanisms of plastid gene expression. *Crit. Rev. Plant Sci.* **12**, 19–55.
- Gruissem, W., and Zurawski, G. (1985). Analysis of promoter regions of the spinach chloroplast *rbcL*, *atpB* and *psbA* genes. *EMBO J.* **4**, 3375–3383.
- Hajdukiewicz, P.T.J., Allison, L., and Maliga, P. (1997). The two RNA polymerase encoded by the nuclear and the plastid compartments transcribe distinct groups of genes in tobacco plastids. *EMBO J.* **16**, 4041–4048.
- Hanley-Bowdoin, L., Orozco, E.M., and Chua, N.-H. (1985). *In vitro* synthesis and processing of a maize chloroplast transcript encoded by the ribulose 1,5-bisphosphate carboxylase large subunit gene. *Mol. Cell. Biol.* **5**, 2733–2745.
- Hayes, R., Kudla, J., Schuster, G., Gabay, L., Maliga, P., and Gruissem, W. (1996). Chloroplast mRNA 3'-end processing by a high molecular weight protein complex is regulated by nuclear encoded RNA binding proteins. *EMBO J.* **15**, 1132–1141.
- Hedtkke, B., Börner, T., and Weihe, A. (1997). Mitochondrial and chloroplast phage-type RNA polymerases in *Arabidopsis*. *Science* **277**, 809–811.
- Hoffer, P.H., and Christopher, D.A. (1997). Structure and blue-light-responsive transcription of a chloroplast *psbD* promoter from *Arabidopsis thaliana*. *Plant Physiol.* **115**, 213–222.
- Hübschmann, T., and Börner, T. (1998). Characterization of transcript initiation sites in ribosome-deficient barley plastids. *Plant Mol. Biol.* **36**, 493–496.
- Igloi, G.L., and Kössel, H. (1992). The transcription apparatus of chloroplasts. *Crit. Rev. Plant Sci.* **10**, 525–558.
- Isono, K., Shimizu, M., Yoshimoto, K., Niwa, Y., Satoh, K., Yokota, A., and Kobayashi, H. (1997a). Leaf-specifically expressed genes for polypeptides destined for chloroplasts with domains for  $\sigma^{70}$  factors of bacterial RNA polymerases in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **94**, 14948–14953.
- Isono, K., Niwa, Y., Satoh, K., and Kobayashi, H. (1997b). Evidence for transcriptional regulation of plastid photosynthesis genes in *Arabidopsis thaliana* roots. *Plant Physiol.* **114**, 623–630.
- Kapoor, S., Suzuki, J.Y., and Sugiura, M. (1997). Identification and functional significance of a new class of non-consensus-type plastid promoters. *Plant J.* **11**, 327–337.
- Kawaguchi, H., Fukuda, I., Shiina, T., and Toyoshima, Y. (1992). Dynamical behavior of *psb* gene transcripts in greening wheat seedlings. I. Time course of accumulation of the *psbA* through *psbN* gene transcripts during light-induced greening. *Plant Mol. Biol.* **20**, 695–704.
- Kim, M., and Mullet, J.E. (1995). Identification of a sequence-specific DNA binding factor required for transcription of the barley chloroplast blue light-responsive *psbD-psbC* promoter. *Plant Cell* **7**, 1445–1457.
- Klein, R.R., and Mullet, J.E. (1990). Light-induced transcription of chloroplast genes: *psbA* transcription is differentially enhanced in illuminated barley. *J. Biol. Chem.* **265**, 1895–1902.
- Klein, U., Salvador, M.L., and Bogorad, L. (1994). Activity of the *Chlamydomonas* chloroplast *rbcL* gene promoter is enhanced by a remote sequence element. *Proc. Natl. Acad. Sci. USA* **91**, 10819–10823.



- Lam, E., Hanley-Bowdoin, L., and Chua, N.-H. (1988). Characterization of a chloroplast sequence-specific DNA binding factor. *J. Biol. Chem.* **263**, 8288–8293.
- Lerbs-Mache, S. (1993). The 110-kDa polypeptide of spinach plastid DNA-dependent RNA polymerase: Single-subunit enzyme or catalytic core of multimeric enzyme complexes? *Proc. Natl. Acad. Sci. USA* **90**, 5509–5513.
- Liu, B., and Troxler, R.F. (1996). Molecular characterization of a positively photoregulated nuclear gene for a chloroplast RNA polymerase  $\sigma$  factor in *Cyanidium caldarium*. *Proc. Natl. Acad. Sci. USA* **93**, 3313–3318.
- Maliga, P. (1998). Two plastid RNA polymerases of higher plants: An evolving story. *Trends Plant Sci.* **3**, 4–6.
- Mayfield, S.P., Yohn, C.B., Cohen, A., and Danon, A. (1995). Regulation of chloroplast gene expression. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **46**, 147–166.
- Mullet, J.E., Orozco, E.M., and Chua, N.-H. (1985). Multiple transcripts for higher plant *rbcL* and *atpB* genes and localization of the transcription initiation sites of the *rbcL* gene. *Plant Mol. Biol.* **4**, 39–54.
- Murashige, T., and Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Plant.* **15**, 473–497.
- Pfannschmidt, T., and Link, G. (1994). Separation of two classes of plastid DNA-dependent RNA polymerases that are differentially expressed in mustard (*Sinapis alba* L.) seedlings. *Plant Mol. Biol.* **25**, 69–81.
- Pfannschmidt, T., and Link, G. (1997). The A and B forms of plastid DNA-dependent RNA polymerase from mustard (*Sinapis alba* L.) transcribe the same genes in a different developmental context. *Mol. Gen. Genet.* **257**, 35–44.
- Reinbothe, S., Reinbothe, C., Heintzen, C., Seidenbecher, C., and Parthier, B. (1993). A methyl jasmonate-induced shift in the length of the 5' untranslated region impairs translation of the plastid *rbcL* transcripts in barley. *EMBO J.* **12**, 1505–1512.
- Rochaix, J.D. (1995). *Chlamydomonas reinhardtii* as the photosynthetic yeast. *Annu. Rev. Genet.* **29**, 209–230.
- Ross, W., Gosink, K.K., Salomon, J., Igarashi, K., Zou, C., Ishihama, A., Severinov, K., and Gourse, R.L. (1993). A third recognition element in bacterial promoters: DNA binding by the  $\alpha$  subunit of the RNA polymerase. *Science* **262**, 1407–1413.
- Sakamoto, W., Kindle, K.L., and Stern, D.B. (1993). *In vivo* analysis of *Chlamydomonas* chloroplast *petD* gene expression using stable transformation of  $\beta$ -glucuronidase translational fusions. *Proc. Natl. Acad. Sci. USA* **90**, 497–501.
- Sakamoto, W., Chen, X., Kindle, K.L., and Stern, D.B. (1994). Function of the *Chlamydomonas reinhardtii* *petD* 5' untranslated region in regulating the accumulation of subunit IV of the cytochrome *b<sub>6</sub>/f* complex. *Plant J.* **6**, 503–512.
- Salvador, M.L., Klein, U., and Bogorad, L. (1993). 5' sequences are important positive and negative determinants of the longevity of *Chlamydomonas* chloroplast gene transcripts. *Proc. Natl. Acad. Sci. USA* **90**, 1556–1560.
- Satoh, J., Baba, K., Nakahira, Y., Shiina, T., and Toyoshima, Y. (1997). Characterization of dynamics of the *psbD* light-induced transcription in mature wheat chloroplasts. *Plant Mol. Biol.* **33**, 267–278.
- Schrubar, H., Wanner, G., and Westhoff, P. (1990). Transcriptional control of plastid gene expression in greening *Sorghum* seedlings. *Planta* **183**, 101–111.
- Shinozaki, K., and Sugiura, M. (1982). The nucleotide sequence of the tobacco chloroplast gene for the large subunit of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Gene* **20**, 91–102.
- Shinozaki, K., Ohme, M., Tanaka, M., Wakasugi, T., Hayashida, N., Matsubayashi, T., Zaita, N., Chunwongse, J., Obokata, J., Yamaguchi-Shinozaki, K., Ohto, C., Torazawa, K., Meng, B.Y., Sugita, M., Deno, H., Kamogashira, T., Yamada, K., Kusuda, J., Takaiwa, F., Kata, A., Todoh, N., Shimada, H., and Sugiura, M. (1986). The complete nucleotide sequence of the tobacco chloroplast genome: Its gene organization and expression. *EMBO J.* **5**, 2043–2049.
- Silhavy, D., and Maliga, P. (1998a). Mapping of promoters for the nucleus-encoded plastid RNA polymerase (NEP) in the *iojap* maize mutant. *Curr. Genet.* **33**, 340–344.
- Silhavy, D., and Maliga, P. (1998b). Plastid promoter utilization in rice embryogenic cell culture. *Curr. Genet.* **34**, 67–70.
- Staub, J.M., and Maliga, P. (1994). Novel extrachromosomal elements in tobacco plastids. *Proc. Natl. Acad. Sci. USA* **91**, 7468–7472.
- Stern, D.B., and Grissem, W. (1987). Control of plastid gene expression: 3' inverted repeats act as mRNA processing and stabilizing elements, but do not terminate transcription. *Cell* **51**, 1145–1157.
- Stern, D.B., and Kindle, K.L. (1993). 3' end maturation of the *Chlamydomonas reinhardtii* chloroplast *atpB* mRNA is a two-step process. *Mol. Cell. Biol.* **13**, 2277–2285.
- Stern, D.B., Radwanski, E.R., and Kindle, K.L. (1991). A 3' stem/loop structure of the *Chlamydomonas* chloroplast *atpB* gene regulates mRNA accumulation in vivo. *Plant Cell* **3**, 285–297.
- Svab, Z., and Maliga, P. (1993). High-frequency plastid transformation in tobacco by selection for a chimeric *aadA* gene. *Proc. Natl. Acad. Sci. USA* **90**, 913–917.
- Svab, Z., Hajdukiewicz, P., and Maliga, P. (1990). Stable transformation of plastids in higher plants. *Proc. Natl. Acad. Sci. USA* **87**, 8526–8530.
- Tanaka, K., Tozawa, Y., Mochizuki, N., Shinozaki, K., Nagatani, A., Wakasa, K., and Takahashi, H. (1997). Characterization of three cDNA species encoding plastid RNA polymerase sigma factors in *Arabidopsis thaliana*: Evidence for the sigma factor heterogeneity in higher plant plastids. *FEBS Lett.* **413**, 309–313.
- Tiller, K., and Link, G. (1993a). Sigma-like transcription factors from mustard (*Sinapis alba* L.) etioplast are similar in size to, but functionally distinct from, their chloroplast counterparts. *Plant Mol. Biol.* **21**, 503–513.
- Tiller, K., and Link, G. (1993b). Phosphorylation and dephosphorylation affect functional characteristics of chloroplast and etioplast transcription systems from mustard (*Sinapis alba* L.). *EMBO J.* **12**, 1745–1753.
- Tiller, K., Eisermann, A., and Link, G. (1991). The chloroplast transcription apparatus from mustard (*Sinapis alba* L.): Evidence for three different transcription factors which resemble bacterial  $\sigma$  factors. *Eur. J. Biochem.* **198**, 93–99.
- To, K.Y., Cheng, M.C., Suen, D.F., Mon, D.P., Chen, L.F.O., and Chen, S.C.G. (1996). Characterization of the light-responsive promoter of rice chloroplast *psbD*-C operon and the sequence-specific DNA binding factor. *Plant Cell Physiol.* **37**, 660–666.

- Tozawa, Y., Tanaka, K., Takahashi, H., and Wakasa, K.** (1998). Nuclear encoding of a plastid sigma factor in rice and its tissue- and light-dependent expression. *Nucleic Acids Res.* **26**, 415–419.
- Vera, A., and Sugiura, M.** (1995). Chloroplast rRNA transcription from structurally different tandem promoters: An additional novel-type promoter. *Curr. Genet.* **27**, 280–284.
- Wada, T., Tunoyama, Y., Shiina, T., and Toyoshima, Y.** (1994). *In vitro* analysis of light-induced transcription in the wheat *psbD/C* gene cluster using plastid extracts from dark-grown and short-term illuminated seedlings. *Plant Physiol.* **104**, 1259–1267.
- Yang, J., Schuster, G., and Stern, D.B.** (1996). CSP41, a sequence-specific chloroplast mRNA binding protein, is an endoribonuclease. *Plant Cell* **8**, 1409–1420.
- Zoubenko, O.V., Allison, L.A., Svab, Z., and Maliga, P.** (1994). Efficient targeting of foreign genes into the tobacco plastid genome. *Nucleic Acids Res.* **22**, 3819–3824.
- Zurawski, G., Whitfield, P.R., and Bottomley, W.** (1986). Sequence of the gene for the large subunit of ribulose-1,5-bisphosphate carboxylase from pea chloroplasts. *Nucleic Acids Res.* **14**, 3975.